

RADIOLOGY ST. PAUL'S HOSPITAL 1081 BURRARD ST VANCOUVER, BC V6Z 1Y6 PH: 604 806-8006 FAX: 604 806-8437

CONSULTATION REPORT

Exam Date: December 15, 2011

cc: 099047 PHYSICIAN DIAGNOSTIC cc: 002841 DEBORAH HOEBERG cc:
 Pt. Name: WALKER, SAMANTHA JUNE

 MPI #: 02174301-8

 DOB:
 41

 Sex: F

 Pt Type: ODOPT
 Loc: PRA

 FRA:

ORD MD: CHEW, JESSE 026618

Report Copies:

E-03508695

CT SCAN OF CHEST

INDICATION: Increased CA 19-9. History of Pseudomyxoma peritonei.

TECHNIQUE: IV contrast-enhanced CT chest.

FINDINGS:

Comparison - CT of the abdomen and pelvis performed December 6, 2011 and December 15, 2010.

There is a 4.2 x 3.8 x 3.6 cm low-density lobulated mass lesion within the anterior segment of the right upper lobe with a small focus of anterior pleural contact measured at 1.7 cm. Additionally there are a number of low density lymph nodes seen within the right hilum measuring up to 1.2 cm in short axis and another low-density lymph node seen more inferiorly just superior to the cavoatrial junction.

Separated from this right upper lobe finding is a lobulated crescentic focus of low density seen in relation to the minor fissure measuring 2.6 x 0.8 cm. As noted on the CT abdomen, there is also a tiny subpleural focus of increased attenuation seen along the posterior border of the right diaphragm unchanged from the most recent prior study. This abnormality was not present on the more distant prior exams.

There is no pericardial or pleural fluid. No aggressive bony lesions are seen.

IMPRESSION:

Lobulated mass lesion within the right upper lobe, associated pleural based findings and right hilar adenopathy. Given the patient's history of prior pseudomyxoma peritonei, these abnormalities would be suspicious for metastatic deposits, even given that parenchymal lung metastasis from this

Exam #: 350869	5 Or	der # 251	9723
Transcribed By:	VPAR	12/16/11	14:44

Dict. Date: 12/15/11 Page 1

Print Date: 12/16/11 Status Date: 12/16/11

PLEASE ENSURE YOUR FAX MACHINE HAS PAPER & IS ON AT ALL TIMES.